

A priori estimates for the Hill and Dirac operators

Evgeny Korotyaev *

February 2, 2008

Abstract

Consider the Hill operator $Ty = -y'' + q'(t)y$ in $L^2(\mathbb{R})$, where $q \in L^2(0, 1)$ is a 1-periodic real potential. The spectrum of T is absolutely continuous and consists of bands separated by gaps $\gamma_n, n \geq 1$ with length $|\gamma_n| \geq 0$. We obtain a priori estimates of the gap lengths, effective masses, action variables for the KDV. For example, if $\mu_n^\pm$ are the effective masses associated with the gap $\gamma_n = (\lambda_n^-, \lambda_n^+)$, then $|\mu_n^- + \mu_n^+| \leq C|\gamma_n|^2 n^{-4}$ for some constant $C = C(q)$ and any $n \geq 1$. In order to prove these results we use the analysis of a conformal mapping corresponding to quasimomentum of the Hill operator. That makes possible to reformulate the problems for the differential operator as the problems of the conformal mapping theory. Then the proof is based on the analysis of the conformal mapping and the identities. Moreover, we obtain the similar estimates for the Dirac operator.

1 Introduction and main results

Consider the self-adjoint periodic weighted operator T_w in $L^2(\mathbb{R}, w^2(t)dt)$, given by

$$T_w f = -w^{-2}(w^2 f')' = -f'' - 2pf', \quad w(t) = e^{\int_0^t p(s)ds}, \quad p \in H = \{p \in L^2(\mathbb{T}), \int_0^1 p(t)dt = 0\}.$$

The operator T_w is well studied, see [K8], [K9], [Kr] and references therein. Define the unitary transformation $U : L^2(\mathbb{R}, w^2 dt) \rightarrow L^2(\mathbb{R}, dt)$ as multiplication by w . Then T_w is unitarily equivalent to the Hill operator $T = UT_w U^{-1}$ given by

$$T = -\left(\frac{d}{dt} + p\right)\left(\frac{d}{dt} - p\right) = -\frac{d^2}{dx^2} + q' + q_0 \geq 0, \quad q' = p'(t) + p^2(t) - \|p\|^2, \quad (1.1)$$

where $q_0 = \|p\|^2 = \int_0^1 p^2(t)dt$ and q' is a 1-periodic potential (distribution). We mention following papers devoted to singular potentials [KM], [K7], [NS], [SS]. We define the Riccati type mapping $R : H \rightarrow H$ by

$$q = R(p) = p(x) + \int_0^x (p^2(t) - \|p\|^2)dt + \int_0^1 \left(t - \frac{1}{2}\right)p^2(t)dt, \quad p \in H. \quad (1.2)$$

*Institut für Mathematik, Humboldt Universität zu Berlin, Rudower Chaussee 25, 12489, Berlin, Germany, e-mail: evgeny@math.hu-berlin.de

Recall that the Riccati map $R : H \rightarrow H$ is a real analytic isomorphism between H and H , see [K7]. Thus, if $p' \in L^2(\mathbb{T})$, then T_w corresponds to the Hill operator T with L^2 -potential.

We can not introduce the standard fundamental solutions for the operator T , since the perturbation q' is very strong. But the operator $T_w = U^{-1}TU$ has the standard fundamental solutions $\varphi(t, \lambda), \vartheta(t, \lambda)$, which satisfy the equation $-y'' - 2py' = \lambda y$, $\lambda \in \mathbb{C}$ and the conditions $\varphi'(0, \lambda) = \vartheta(0, \lambda) = 1$, $\varphi(0, \lambda) = \vartheta'(0, \lambda) = 0$. Here and below we use the notation $u' = \frac{\partial u}{\partial t}$. Introduce the Lyapunov function $\Delta(\lambda) = \frac{\varphi'(1, \lambda) + \vartheta(1, \lambda)}{2}$.

It is well known, that the spectrum $\sigma(T) = \sigma(T_w)$ of T is absolutely continuous and consists of intervals $\sigma_n = [\lambda_{n-1}^+, \lambda_n^-]$, where $\lambda_{n-1}^+ < \lambda_n^- \leq \lambda_n^+$, $n \geq 1$ and $\lambda_0^+ = 0$. The intervals σ_n and σ_{n+1} are separated by the gap $\gamma_n = (\lambda_n^-, \lambda_n^+)$. If a gap degenerates, that is $\gamma_n = \emptyset$, then the corresponding segments σ_n and σ_{n+1} merge. Note that $\Delta(\lambda_n^\pm) = (-1)^n$, $n \geq 1$, and there exists a unique point $\lambda_n \in [\lambda_n^-, \lambda_n^+]$ such that $\Delta'(\lambda_n) = 0$. The sequence $\lambda_0^+ < \lambda_1^- \leq \lambda_1^+ < \dots$ is the spectrum of equation $-y'' + qy = zy$ with the 2-periodic boundary conditions, i.e. $y(t+2) = y(t)$, $t \in \mathbb{R}$. If $\lambda_n^- = \lambda_n^+$ for some n , then this number $\lambda_n^\pm$ is the double eigenvalue of this equation with the 2-periodic boundary conditions. The lowest eigenvalue λ_0^+ is always simple, $\Delta(\lambda_0^+) = 1$, and the corresponding eigenfunction is 1-periodic. The eigenfunctions, corresponding to the eigenvalue $\lambda_n^\pm$, are 1-periodic, when n is even and they are antiperiodic, i.e. $y(t+1) = -y(t)$, $t \in \mathbb{R}$, when n is odd.

We introduce a new parameter (momentum) $z = x + iy = \sqrt{\lambda}$, $\sqrt{1} = 1$, and momentum gaps

$$g_n = (z_n^-, z_n^+), \quad z_n^\pm = \sqrt{\lambda_n^\pm} \geq 0, \quad g_{-n} = -g_n, \quad n \geq 1, \quad g_0 = \emptyset.$$

We define a quasimomentum $k(z) = u(z) + iv(z) = \arccos \Delta(z^2)$, $z \in \mathcal{Z} = \mathbb{C} \setminus \overline{g}$, $g = \cup g_n$, see [MO]. The function $k(z)$ is a conformal mapping from $\mathcal{Z}$ onto a quasimomentum domain $\mathcal{K} = \mathbb{C} \setminus \cup_{n \in \mathbb{Z}} \Gamma_n$, where the slit $\Gamma_n = [\pi n + ih_n, \pi n - ih_n]$, $h_0 = 0$, and $h_n = h_{-n} \geq 0$ is defined by the equation $\Delta(\lambda_n) = (-1)^n \cosh h_n$. The quasimomentum k satisfies $k(z) = z + o(1)$ as $|z| \rightarrow \infty$. With each edge of the gap $\gamma_n \neq \emptyset$, we associate the effective mass $\mu_0^+, \mu_n^\pm$ by

$$\lambda = z^2 = \lambda_n^\pm + \frac{(k - \pi n)^2}{2\mu_n^\pm}(1 + o(1)) \quad \text{as} \quad \lambda \rightarrow \lambda_n^\pm. \quad (1.3)$$

Introduce the real Hilbert spaces $\ell_p^2 = \left\{ f = (f_n)_1^\infty, \quad f_n \in \mathbb{R}, \quad \|f\|_p^2 = \sum_{n \geq 1} (2\pi n)^{2p} f_n^2 < \infty \right\}$, $p \geq 0$ and let $\ell_0^2 \equiv \ell^2$. For the sake of the reader, we briefly recall the results existing

in the literature about the a priori estimates. Marchenko and Ostrovski [MO1-2] obtained the estimates: $\|q_1\| \leq C(1 + h_+) \|h\|_1$, $\|h\|_1 \leq C\|q_1\| \exp(A\|q_1\|)$, for some absolute constants A, C , where $h_+ = \sup_{n \geq 1} h_n$ and $q_1 = q_0 + q'$. These estimates are not sharp since they used the Bernstein inequality. Using the harmonic measure argument Garnett and Trubowitz [GT] obtained $\|\gamma\| \leq (4 + \|h\|_1) \|h\|_1$.

Various estimates for the Hill operator with a potential $q' \in L^2(0, 1)$ were obtained in the author's paper [K5-6] and in [KK1], [K3] for the Dirac operator. In the paper [K3], [K5], [K6] the following sharp estimates for the case $q' \in L^2(0, 1)$ were obtained:

$$\|\gamma\| \leq 6\|q'\|(1 + \|q'\|)^{\frac{1}{3}}, \quad \|q'\| \leq 4\|\gamma\|(1 + \|\gamma\|)^{\frac{1}{3}}, \quad \gamma = (|\gamma_n|)_1^\infty,$$

$$2\|h\|_1 \leq \pi\|q'\|(1 + \|q'\|)^{\frac{1}{3}}, \quad \|q'\| \leq 3\|h\|_1(6 + h_+)^{\frac{1}{2}}, \quad h_+ = \sup_{n \geq 1} h_n,$$

and for the case $q \in L^2(0, 1)$ [K8]:

$$\|q\| \leq 2\|h\|(1 + 4\|h\|), \quad \|h\| \leq 3\|q\|(1 + 2\|q\|), \quad q \in H,$$

$$\|q\| \leq 48\pi^2\|\gamma\|_{-1}(1 + \|\gamma\|_{-1})^3, \quad \|\gamma\|_{-1} \leq 2\|q\|(1 + 2\|q\|)^3.$$

It is well known that $v(z) = \text{Im } k(z) \geq v_n(z) = |(z - z_n^-)(z - z_n^+)|^{\frac{1}{2}}, z \in g_n$, see [KK1]. Define a function $V_n(z) = \frac{v(z)}{v_n(z)} - 1 \geq 0, z \in g_n$, which is analytic in the domain $\mathbb{C}_+ \cup \mathbb{C}_- \cup (z_{n-1}^+, z_{n+1}^-)$, see [KK1]. Let $M_n = \max_{z \in g_n} V_n(z), M'_n = \max_{z \in g_n} V'_n(z), \dots$

Theorem 1.1. *Let real $q \in L^2(0, 1)$. Then for each $n \geq 1$ the following estimates hold*

$$|\lambda_n^0 - \frac{|g_n|^2}{4} - \lambda_n| \leq \frac{3|\gamma_n|^2}{8z_n^0} M'_n, \quad \text{where } z_n^0 = \frac{z_n^- + z_n^+}{2}, \quad \lambda_n^0 = \frac{\lambda_n^- + \lambda_n^+}{2}, \quad (1.4)$$

$$|h_n - \frac{|\gamma_n|(1 + V_n(z_n))}{4z_n^0}| \leq \frac{|\gamma_n|^3 M_n'^2}{16z_n^0{}^3}, \quad (1.5)$$

$$|h_n - h_n^\pm| \leq \frac{|g_n|^2}{2} M'_n \left(1 + \frac{M'_n \sqrt{|\gamma_n| |\mu_n^\pm|}}{4} \right), \quad \text{where } h_n^\pm = \left| |g_n| z_n^\pm \mu_n^\pm \right|^{\frac{1}{2}}, \quad (1.6)$$

$$|\mu_n^+ + \mu_n^-| \leq \frac{|g_n|^2}{4z_n^+ z_n^-} (1 + M_n)(1 + 2z_n^0 M'_n), \quad (1.7)$$

$$|\mu_n^\pm \mp \mu_n| \leq \frac{|g_n|^2}{8z_n^0 z_n^\pm} (1 + M_n)^2 \left[1 + 4z_n^\pm M'_n \left(1 + \frac{M'_n}{4} \sqrt{|\gamma_n| |\mu_n^\pm|} \right) \right], \quad \text{where } \mu_n = \frac{2h_n^2}{|\gamma_n|}. \quad (1.8)$$

Remark. 1) Using Theorem 1.1 and estimates

$$|\gamma_n| \leq (4\pi n)^2 (\mu_n^+ - \mu_n^-), \quad h_n^2 \leq \frac{(3\pi)^2}{2} |\gamma_n| |\mu_n^\pm|, \quad (1.9)$$

from [K4], we obtain the estimates in terms of effective masses.

2) Recall that $z_n^\pm = \pi n \pm |\int_0^1 e^{i2\pi n x} q(x) dx| + \ell^d(n)$, $d > 1$ as $n \rightarrow \infty$, see [K8]. This yields asymptotic estimate $\rho_n \geq \pi - \varepsilon$ as $n \rightarrow \infty$, for any $\varepsilon > 0$.

Define the action variables $A_n = \frac{4}{\pi} \int_{g_n} x v(x) dx \geq 0$, $n \geq 1$ for the KDV equation [FM].

Theorem 1.2. *Let real $q \in L^2(0, 1)$ and let $\beta_n = \sqrt{A_n} \geq 0, n \geq 1$. Then the following estimates hold*

$$\left| A_n - \frac{|\gamma_n|^2}{8z_n^0} (1 + V_n(z_n^0)) \right| \leq \frac{|g_n|^4}{2^5} (z_n^0 M_n'' + M'_n), \quad (1.10)$$

$$|A_n - \frac{|\gamma_n| h_n}{2}| \leq \frac{|g_n|^4}{2^5} (z_n^0 M_n'' + 6z_n^0 M_n'^2 + M'_n), \quad (1.11)$$

$$|A_n - \frac{|\gamma_n|}{2} \left| \frac{|g_n| m_n^\pm}{2} \right|^{\frac{1}{2}}| \leq \frac{|g_n|^3}{4} \left(z_n^0 M'_n + \frac{|g_n|}{8} M'_n + |g_n| M_n'' \right), \quad (1.12)$$

$$\left| \beta_n - |\gamma_n| \sqrt{\frac{(1 + V_n(z_n^0))}{8z_n^0}} \right| \leq \frac{|g_n|^3}{2^5 \sqrt{2z_n^0}} (z_n^0 M_n'' + M_n'), \quad (1.13)$$

$$\left| \beta_n - \sqrt{\frac{|\gamma_n| h_n}{2}} \right| \leq \frac{|g_n|^3}{2^5 \sqrt{2z_n^0}} (z_n^0 M_n'' + 6z_n^0 M_n'^2 + M_n'), \quad (1.14)$$

$$|\beta_n - \sqrt{\frac{|\gamma_n|}{2}} \frac{|g_n| m_n^\pm|^{\frac{1}{4}}}{2^{\frac{1}{4}}}| \leq \frac{|g_n|^2}{4 \sqrt{2z_n^0}} \left(z_n^0 M_n' + \frac{|g_n|}{8} M_n' + |g_n| M_n'' \right). \quad (1.15)$$

Consider the Dirac operator T_D **acting in the space** $L^2(\mathbb{R}) \oplus L^2(\mathbb{R})$ **and given by**

$$T_D = \mathcal{J} \frac{d}{dt} + q_D(t), \quad q_D = \begin{pmatrix} q_1 & q_2 \\ q_2 & -q_1 \end{pmatrix} \quad \mathcal{J} \equiv \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},$$

where $q_1, q_2 \in L^2(0, 1)$ are real 1-periodic functions of $t \in \mathbb{R}$. The spectrum of T_D is purely absolutely continuous and $\sigma(T_D) = \cup_{n \in \mathbb{Z}} s_n$, where $s_n = [z_{n-1}^+, z_n^-]$ are spectral bands and $z_{n-1}^+ < z_n^- \leq z_n^+$, (see [LS] for details). The intervals s_n, s_{n+1} are separated by the gap $g_n = (z_n^-, z_n^+)$ with length $|g_n| \geq 0$. If a gap $g_n = \emptyset$, then the corresponding spectral bands s_n, s_{n+1} merge. Introduce the 2×2 -matrix valued fundamental solution $\psi = \psi(t, z)$ satisfying $J\psi' + q_D\psi = z\psi$, $\psi(0, z) = I_2$, $z \in \mathbb{C}$, where I_2 is the 2×2 identity matrix. Define the Lyapunov function $\Delta(z) = \frac{1}{2} \text{Tr} \psi(1, z)$. Note that $\Delta(z_n^\pm) = (-1)^n$, $n \in \mathbb{Z}$, and the function $\Delta'(z)$ has a zero z_n in a "closed gap" $[z_n^-, z_n^+]$ (see e.g. [LS]). The sequence $..z_0^+ < z_1^- \leq z_1^+ < \dots$ is the spectrum of equation $Jy' + q_D y = zy$, $y = (y_1, y_2)^\top$, with the 2-periodic boundary conditions, i.e. $y(t+2) = y(t)$, $t \in \mathbb{R}$. If $z_n^- = z_n^+$ for some n , then $z_n^\pm$ is the double eigenvalue of the equation $Jy' + q_{zs} y = zy$ with the 2-periodic boundary conditions. The eigenfunctions, corresponding to the eigenvalue $z_n^\pm$, are 1-periodic, when n is even and they are antiperiodic, i.e. $y(t+1) = -y(t)$, $t \in \mathbb{R}$, when n is odd.

For each q_D there exists a unique conformal mapping (the quasimomentum) $k : \mathcal{Z} \rightarrow \mathcal{K} = \mathbb{C} \setminus \cup \Gamma_n$, $\Gamma_n = [\pi n - ih_n, \pi n + ih_n]$, given by (see [K6])

$$\cos k(z) = \Delta(z), \quad z \in \mathcal{Z} = \mathbb{C} \setminus \bar{g}, \quad g = \cup g_n, \quad k(z) = z + o(1) \text{ as } |z| \rightarrow \infty,$$

where the height $h_n \geq 0$ is defined by the equation $\cosh h_n = (-1)^n \Delta(z_n) \geq 1$. With each edge of the gap $g_n \neq \emptyset$, we associate the effective mass $m_n^\pm$ by

$$z = z_n^\pm + \frac{(k(z) - \pi n)^2}{2m_n^\pm} (1 + o(1)) \quad \text{as } z \rightarrow z_n^\pm. \quad (1.16)$$

Theorem 1.3. *Let real $q_D \in L^2(0, 1)$. Then for each $n \in \mathbb{Z}$ the following estimates hold*

$$|z_n - z_n^0| \leq \frac{|g_n|^2}{4} M_n', \quad (1.17)$$

$$|2h_n - |g_n|(1 + V_n(z_n))| \leq \frac{|g_n|^3}{8} M_n'^2, \quad (1.18)$$

$$|h_n - h_n^\pm| \leq \frac{|g_n|^2}{2} M'_n \left(1 + \frac{M'_n}{8} \sqrt{2|g_n||m_n^\pm|} \right), \quad \text{where } h_n^\pm = \sqrt{\frac{|g_n||m_n^\pm|}{2}}, \quad (1.19)$$

$$|m_n^+ + m_n^-| \leq |g_n|^2 (1 + M_n) M'_n, \quad (1.20)$$

$$|m_n^\pm \mp m_n| \leq |g_n|^2 (1 + M_n) M'_n \left(1 + \frac{M'_n}{8} \sqrt{2|g_n||m_n^\pm|} \right), \quad \text{where } m_n = \frac{2h_n^2}{|g_n|}. \quad (1.21)$$

Remark. Using Theorem 1.3 and estimates

$$\frac{|g_n|}{2} \leq h_n \leq \pi \sqrt{2|g_n||m_n^\pm|} \leq 2\pi |m_n^\pm|, \quad h_n^2 \leq 2|g_n| \sqrt{m_n^+ m_n^-}, \quad |g_n| \leq 2|m_n^\pm|. \quad (1.22)$$

from [KK1], we obtain the estimates in terms of effective masses.

Recall that the action variables for the NLS equation are given by $a_n = \frac{1}{\pi} \int_{g_n} v(x) dx$, $n \in \mathbb{Z}$.

Theorem 1.4. *Let real $q_D \in L^2(0, 1)$ and let $b_n = \sqrt{a_n} \geq 0, n \in \mathbb{Z}$. Then the following estimates hold*

$$|a_n - (|g_n|^2/8)(1 + V_n(z_n^0))| \leq 2 \frac{|g_n|^4}{4^4} M''_n, \quad (1.23)$$

$$|a_n - \frac{|g_n|h_n}{4}| \leq \frac{|g_n|^4}{2^7} (M''_n + 6M_n'^2), \quad (1.24)$$

$$|a_n - \frac{|g_n|^{3/2}}{2^{5/2}} |m_n^\pm|^{1/2}| \leq \frac{|g_n|^3}{16} M'_n + \frac{|g_n|^4}{2^7} M''_n, \quad (1.25)$$

$$|b_n - 2^{-3/2} |g_n| \sqrt{1 + V_n(z_n^0)}| \leq \frac{\sqrt{2}|g_n|^3}{4^3} M''_n, \quad (1.26)$$

$$|b_n - \sqrt{|g_n|h_n/2}| \leq \frac{\sqrt{2}|g_n|^3}{2^7} (M''_n + 6M_n'^2), \quad (1.27)$$

$$|b_n - \frac{|g_n|^{3/4}}{2^{5/4}} |m_n^\pm|^{1/4}| \leq \frac{(\frac{|g_n|^2}{8} M'_n + \frac{|g_n|^3}{8^2} M''_n)}{\sqrt{2}}. \quad (1.28)$$

Note that Theorem 1.4 is used to study electrostatic inverse problems on the plane [KK3].

Define integrals $Q_p = \frac{1}{\pi} \int_{\mathbb{R}} x^p v(x) dx, p \in \mathbb{Z}$. Recall the identities for the Hill operator

$$Q_0 = \frac{q_0}{2} = \frac{\|p\|^2}{2}, \quad Q_2 = \frac{1}{8} \int_0^1 (q_0^2 + q'(t)^2) dt, \quad (1.29)$$

(recall that $\lambda_0^+ = 0$ for the Hill operator), and for the Dirac operator

$$Q_0 = \frac{\|q_D\|^2}{2} = \frac{1}{2} \int_0^1 (q_1^2(t) + q_2^2(t)) dt, \quad Q_2 = \frac{1}{8} \int_0^1 ((q_1'^2 + q_2'^2) + (q_1^2 + q_2^2)^2) dt. \quad (1.30)$$

Theorem 1.5. *Let real $q \in L^2(0, 1)$ or $q_D \in L^2(0, 1)$. Let $\rho_n = \text{dist}(g_n, g \setminus g_n)$, $n \in \mathbb{Z}$. Then for each $n \in \mathbb{Z}$ the following estimates hold*

$$M_n \leq \frac{2}{\pi \rho_n} \sup_{s \in \mathbb{Z}} h_s \leq \frac{\sqrt{8Q_0}}{\pi \rho_n}, \quad M'_n \leq \frac{M_n}{\rho_n}, \quad M''_n \leq \frac{M_n}{\rho_n^2}, \quad (1.31)$$

$$\sum_{n \in \mathbb{Z}} M_n \leq \frac{\pi^2}{3\rho^2} Q_0, \quad \text{where} \quad \rho = \inf_{n \in \mathbb{Z}} \rho_n, \quad (1.32)$$

$$\frac{1}{\rho} \leq \frac{5}{2} e^{5\sqrt{Q_0/2}}. \quad (1.33)$$

If $q' \in L^2(0, 1)$ or $q'_D \in L^2(0, 1)$, then

$$V_n(x) \leq \frac{2}{|x| \sqrt{|z_n^+ z_n^-|}} \left(Q_0 + \frac{Q_2}{\rho_n^2} \right), \quad x \in \bar{g}_n. \quad (1.34)$$

Corollary 1.6. *Let real $q_D \in L^2(0, 1)$. Then the following estimates hold*

$$\left(\sum_{n \in \mathbb{Z}} |a_n - \frac{|g_n| h_n}{4}|^{\frac{1}{3}} \right)^3 \leq Q_0^3 \left(\frac{\pi^2}{4^7 \rho^4} \right) \left(1 + \frac{\sqrt{8Q_0}}{\pi \rho} \right), \quad (1.35)$$

$$\sum_{n \in \mathbb{Z}} |a_n - \frac{|g_n| h_n}{4}| \leq \frac{g_+^4}{2^7} \frac{\pi^2 Q_0}{3\rho^4} \left(1 + 6 \frac{\sqrt{8Q_0}}{\pi \rho} \right), \quad g_+ = \sup_n |g_n|. \quad (1.36)$$

In order to prove these results we consider the quasimomentum of the Hill (or the Dirac) operator as a conformal mapping. That makes possible to reformulate the problem for the differential operator as a problem of the conformal mapping theory (see [KK1], [K1]-[K8] and [MO1]). We use the Poisson integral for the domain $\mathbb{C}_+ \cup \mathbb{C}_- \cup g_n$ and some a priori estimates from [KK1]. The results of this paper are used in [KK3].

2 Estimates for conformal mappings

Consider a conformal mapping $z : \mathcal{K}_+ \rightarrow \mathbb{C}_+$ with asymptotics $z(iv) = iv(1 + o(1))$ as $v \rightarrow \infty$, where $z(k), k = u + iv \in \mathbb{K}$. The domain $\mathcal{K}_+ = \mathbb{C}_+ \cap \mathcal{K}$ for some sequence $(h_n)_{n \in \mathbb{Z}} \in \ell^\infty, h_n \geq 0$ and the domain $\mathcal{K} = \mathbb{C} \setminus \cup_{n \in \mathbb{Z}} \Gamma_n$, where $\Gamma_n = [u_n - ih_n, u_n + ih_n]$ and $u_n, n \in \mathbb{Z}$ is strongly increasing sequence of real numbers such that $u_n \rightarrow \pm\infty$ as $n \rightarrow \pm\infty$. The difference of any two such mappings equals a real constant. Thus the imaginary part $y(k) = \text{Im } z(k)$ is unique. We call such mapping $z(k)$ the comb mapping. Define the inverse mapping $k(\cdot) : \mathbb{C}_+ \rightarrow \mathcal{K}_+$. It is clear that $k(z), z = x + iy \in \mathbb{C}_+$ has the continuous extension into $\bar{\mathbb{C}}_+$. We define "gaps" g_n , "bands" s_n and the "spectrum" s of the comb mapping by:

$$g_n = (z_n^-, z_n^+) = (z(u_n - 0), z(u_n + 0)), \quad s_n = [z_{n-1}^+, z_n^-], \quad s = \cup_{n \in \mathbb{Z}} s_n.$$

It is well known that the set s can not be the spectrum of two different comb mappings [Le]. The function $u(z) = \text{Re } k(z)$ is strongly increasing on each band s_n and $u(z) = u_n$ for all

$z \in [z_n^-, z_n^+]$, $n \in \mathbb{Z}$; the function $v(z) = \text{Im } k(z)$ equals zero on each band s_n and is strongly convex on each gap $g_n \neq \emptyset$ and has the maximum at some point z_n given by $v(z_n) = h_n$. If the gap is empty we set $z_n = z_n^\pm$. The function $z(\cdot)$ has an analytic extension (by the symmetry) from the domain $\mathcal{K}_+$ onto the domain $\mathcal{K}$ and $z(\cdot) : \mathcal{K} \rightarrow z(\mathcal{K}) = \mathcal{Z} = \mathbb{C} \setminus \bigcup \bar{g}_n$ is a conformal mapping. For each gap $g_n = (z_n^-, z_n^+) \neq \emptyset$ we define the effective masses $m_n^\pm$ by

$$z(k) - z_n^\pm = \frac{(k - u_n)^2}{2m_n^\pm} + O((k - u_n)^3) \quad \text{as } z \rightarrow z_n^\pm. \quad (2.1)$$

If $|g_n| = 0$, then we set $m_n^\pm = 0$. It is clear that if $|g_n| > 0$, then $\pm m_n^\pm > 0$. These and others properties of the comb mappings it is possible to find in the papers of Levin [Le].

We emphasize that the comb mapping $k(z)$ with $u_n = \pi n, n \in \mathbb{Z}$ corresponds to the quasimomentum for the Dirac operator with periodic coefficients. If the comb mapping $k(z)$ is symmetric, $k(-z) = -k(z), z \in \mathcal{Z}$, with $u_n = \pi n, n \in \mathbb{Z}$, then it corresponds to the quasimomentum for the Hill operator. Furthermore, the comb mapping is the rotation number for the Schrödinger operator with some almost periodic potential (see [JM]).

For some $(h_n)_{n \in \mathbb{Z}} \in \ell_\mathbb{R}^\infty$ we fix some gap $G = (-c, c), c > 0$. The function $v = \text{Im } k(z)$ satisfies

$$v(x) = r(x)(1 + V(x)), \quad V(x) = \frac{1}{\pi} \int_{\mathbb{R} \setminus G} \frac{v(t) dt}{|t - x| r(t)}, \quad r(x) = |c^2 - x^2|^{\frac{1}{2}}, \quad x \in (-c, c), \quad (2.2)$$

see [KK1]. Recall that $v''(x) < 0$ for all $x \in G$. The function $v(x), x \in G$ has a unique maximum h at some point $z_0 \in (-c, c)$ given by

$$h = v(z_0) = r(z_0)(1 + V_0) \geq c, \quad V_0 = V(z_0). \quad (2.3)$$

Lemma 2.1. *Let a conformal mapping $k : \mathcal{Z} \rightarrow \mathcal{K}$ have a gap $G = (-c, c), c > 0$ for some $(h_n)_{n \in \mathbb{Z}} \in \ell^\infty$ and $V_0 = V(z_0), V'_0 = V'(z_0), \dots, r_0 = r(z_0), \nu = -1/v''(z_0) > 0$. Then*

$$z_0 = \frac{r_0^2 V'_0}{1 + V_0}, \quad \text{where } r_0 \leq c, \quad (2.4)$$

$$|c - r_0| \leq \frac{r_0^3 V_0'^2}{2(1 + V_0)^2}, \quad (2.5)$$

$$|h - c(1 + V_0)| \leq \frac{r_0^3 V_0'^2}{2(1 + V_0)}, \quad (2.6)$$

$$\left| \frac{c^2}{h} - \nu \right| \leq hc^2((3 + c)V_0'^2 + V_0''). \quad (2.7)$$

Proof. We have $v'(z_0) = 0$ and $0 = v'(x) = r'(x)(1 + V(x)) + r(x)V'(x)$ at $x = z_0$ and $(r^2)' = -2x$, which yield (2.4). Substituting (2.4) into the identity $z_0^2 = c^2 - r_0^2$, where $r_0 = r(z_0)$ we obtain $c^2 - r_0^2 = \frac{r_0^4 V_0'^2}{(1 + V_0)^2}$, which implies (2.5).

Estimate (2.5) together with the identity $h = r_0(1 + V_0)$ yield

$$|h - c(1 + V_0)| = (1 + V_0)|r_0 - c| \leq \frac{r_0^3 V_0'^2}{2(1 + V_0)},$$

which gives (2.6). Differentiating the function $v'(x)r(x)$ at z_0 we obtain

$$r_0 v''(z_0) = (rv')'|_{z=z_0} = \left(-x(1 + V) + r^2 V' \right)'|_{z=z_0} = -(1 + V_0) - 3z_0 V_0' + r_0^2 V_0'',$$

which gives $r_0 v''(z_0) = -(1 + V_0) - 3z_0 V_0' + r_0^2 V_0''$. Then the definition $\nu_n = -1/v''(z_0)$ implies $r_0 = \nu(1 + V_0 + 3V_0'z_0 - r_0^2 V_0'')$, which yields

$$\frac{r_0}{1 + V_0} = \frac{r_0^2}{h} = \nu + R, \quad R = \nu \frac{3V_0'z_0 - r_0^2 V_0''}{1 + V_0}, \quad c^2 - r_0^2 = \frac{r_0^4 V_0'^2}{(1 + V_0)^2}. \quad (2.8)$$

Thus $\frac{c^2}{h} = \nu + R_1$, $R_1 = R + \frac{c^2 - r_0^2}{h}$. Thus $|R| \leq \nu r_0^2 (3V_0'^2 + V_0'')$ and $\frac{c^2 - r_0^2}{h} \leq \frac{r_0^3 V_0'^2}{(1 + V_0)^2}$, which yield (2.7), where the estimate $\nu \leq h$ from [KK4] was used. ■

Recall the identity from [KK1]

$$\pm m^\pm = c(1 + V(\pm c)^2) \geq c. \quad (2.9)$$

We obtain estimates of the action. Let $M = \max_{x \in G} V(x)$, $M' = \max_{x \in G} V'(x)$, ...

Lemma 2.2. *Let a conformal mapping $k : \mathcal{Z} \rightarrow \mathcal{K}$ have a gap $G = (-c, c)$, $c > 0$ for some $(h_n)_{n \in \mathbb{Z}} \in \ell^\infty$. Then the actions $a = \frac{1}{\pi} \int_G v(x) dx$ and $A = \frac{4}{\pi} \int_G x v(x) dx$ and $b = |a|^{\frac{1}{2}} > 0$ satisfy*

$$|a - \frac{c^2}{2}(1 + V(0))| \leq \frac{c^4}{8} M'', \quad (2.10)$$

$$|A| \leq \frac{c^4}{2} M', \quad (2.11)$$

$$|a - \frac{ch}{2}| \leq \frac{c^4}{8} (M'' + 6M'^2), \quad (2.12)$$

$$|a - \frac{c}{2} \sqrt{c|m_\pm|}| \leq \frac{c^3}{2} (M' + \frac{c}{4} M''), \quad (2.13)$$

$$|b - c \sqrt{(1 + V(0))/2}| \leq \frac{c^3}{8\sqrt{2}} M'', \quad (2.14)$$

$$|b - \sqrt{ch/2}| \leq \frac{c^3}{8\sqrt{2}} (M'' + 6M'^2), \quad (2.15)$$

$$|b - \frac{c^{3/4}}{\sqrt{2}} |m_\pm|^{\frac{1}{4}}| \leq \frac{c^2}{2\sqrt{2}} (M' + \frac{c}{4} M''). \quad (2.16)$$

Proof. Using the Taylor formula $V(x) = V(0) + V'(0)x + \int_0^x V''(s)(x-s)ds$, we obtain

$$\begin{aligned}\pi a &= \int_G v(x)dx = \int_G r(x)(1+V(x))dx = a_{(1)} + a_{(2)}, \quad a_{(2)} = \int_G r(x)dx \int_0^x V''(s)(x-s)ds, \\ a_{(1)} &= \int_G r(x)(1+V(0)+V'(0)x)dx = \pi \frac{c^2}{2}(1+V(0)).\end{aligned}$$

Consider $a_{(2)}$. The function $V''(x) > 0$ for all $x \in G$, then for some $x_1 \in G$ we get

$$a_{(2)} = \int_G r(x)dx \int_0^x V''(s)(x-s)ds = V''(x_1) \int_G r(x)dx \int_0^x (x-s)ds = \pi V''(x_1) \frac{c^4}{24}$$

which yields (2.10). Using $V(x) = V(0) + \int_0^x V'(s)ds$ and $\int_G xr(x)(1+V(0))dx = 0$, we have

$$\begin{aligned}A &= \frac{4}{\pi} \int_G xv(x)dx = \frac{4}{\pi} \int_G xr(x)dx \int_0^x V'(s)ds, \\ |A| &\leq \frac{4}{\pi} \int_G |x|r(x) \left| \int_0^x V'(s)ds \right| dx \leq \frac{4M'}{\pi} \int_G x^2 r(x)dx = \frac{c^4 M'}{2},\end{aligned}$$

which yields (2.11). Using (2.3) we have the following identity

$$h = c(1+V(0)) + (h - c(1+V_0)) + cz_0 V'(x_1), \quad \text{where } V_0 = V(z_0),$$

for some $x_1 \in G$, which yields

$$a - \frac{hc}{2} = \left(a - \frac{c^2}{2}(1+V(0)) \right) + \frac{c(h - c(1+V_0))}{2} + \frac{c^2 z_0}{2} V'(x_1).$$

Using (2.10), (2.6), (2.4) we obtain

$$\left| a - \frac{hc}{2} \right| \leq \frac{c^4}{2^3} M'' + \frac{c^4 M'^2}{4} + \frac{c^4 M'^2}{2} < \frac{c^4}{8} (M'' + 6M'^2),$$

which yields (2.12). Using (2.9) we obtain

$$\left| a - \frac{c}{2} \sqrt{cm_{\pm}} \right| \leq |a - (c^2/2)(1+V(0))| + \frac{c^2}{2} |V(0) - V(\pm c)| \leq \frac{c^4}{8} M'' + \frac{c^3}{2} M',$$

which yields (2.13). Using (2.10) we obtain $|b - c\sqrt{(1+V(0))/2}| \leq \frac{c^4}{8} \frac{M''}{|b+2^{-\frac{1}{2}}c|}$, and the estimate $a = b^2 \geq c^2/2$ yields (2.14). Using (2.12) we obtain $|b - \sqrt{ch/2}| \leq \frac{c^4}{8} \frac{M''+6M'^2}{|b+\sqrt{ch/2}|}$, and the estimate $a = b^2 \geq c^2/2, h \geq c$ yields (2.15). Due to (2.13) we have

$$\left| b - \frac{c^{3/4}}{\sqrt{2}} |m_{\pm}|^{\frac{1}{4}} \right| \leq \frac{c^3}{2} \frac{(M' + \frac{c}{4} M'')}{(b + \frac{c^{3/4}}{\sqrt{2}} |m_{\pm}|^{\frac{1}{4}})},$$

and $|m_{\pm}| \geq c, a \geq \frac{c^2}{2}$ gives (2.16). ■

Lemma 2.3. *Let a conformal mapping $k : \mathcal{Z} \rightarrow \mathcal{K}$ have a gap $G = (-c, c)$, $c > 0$ for some $(h_n)_{n \in \mathbb{Z}} \in \ell^\infty$. Then*

$$|h - h^\pm| \leq 2c^2 \left(M' + \frac{V_0'^2 \sqrt{cm^\pm}}{4(1+V_0)^2} \right), \quad \text{where } h^\pm = \sqrt{c|m^\pm|}, \quad (2.17)$$

$$|m^+ + m^-| \leq 4c^2(1+M)M', \quad (2.18)$$

$$|m^\pm \mp m| \leq 4c^2(1+M) \left(M' + \frac{V_0'^2 \sqrt{cm^\pm}}{4(1+V_0)^2} \right), \quad \text{where } m = \frac{h^2}{c}. \quad (2.19)$$

Proof. Let $V_0 = V(z_0)$ and $V_\pm = V(\pm c)$. Identities (2.9), (2.3) give

$$|h - h^\pm| = |r_0(1+V_0) - c(1+V_\pm)| \leq r_0|V_0 - V_\pm| + |r_0 - c|(1+V_\pm).$$

Then estimates (2.5), (2.9) give $|r_0 - c|(1+V_\pm) \leq \frac{c^2 V_0'^2 \sqrt{c|m^\pm|}}{2(1+V_0)^2}$, and we also have $|V(z_0) - V(\pm c)| \leq 2cM'$. Combining two last estimates we obtain (2.17).

We prove (2.18). Identity (2.9) gives

$$|m^+ + m^-| = c(2 + V_+ + V_-)|V_+ - V_-| \leq 4c^2(1+M)M',$$

which yields (2.18). We prove (2.19). Using (2.9), (2.3) we have

$$|m^+ - m| = \frac{|(h^+)^2 - h^2|}{c} = \frac{|h^+ - h||h^+ + h|}{c} \leq |h^+ + h|2c \left(M' + \frac{V_0'^2 \sqrt{cm^\pm}}{4(1+V_0)^2} \right),$$

and $h + h^+ \leq 2c(1+M)$ yields (2.19). ■

We rewrite (2.2) for any gap $g_n = (z_n^-, z_n^+)$ by

$$v(x) = v_n(x)(1+V_n(x)), \quad V_n(x) = \frac{1}{\pi} \int_{g \setminus g_n} \frac{v(t)dt}{v_n(t)|t-x|}, \quad v_n(x) = |(x-z_n^-)(x-z_n^+)|^{\frac{1}{2}}, \quad (2.20)$$

$x \in g_n$. We have the following identities and estimates

$$V_n'(x) = \frac{1}{\pi} \int_{g \setminus g_n} \frac{v(t)dt}{(t-x)|t-x|v_n(t)}, \quad V_n''(x) = \frac{2}{\pi} \int_{g \setminus g_n} \frac{v(t)dt}{|t-x|^3 v_n(t)} > 0, \quad (2.21)$$

$$|V_n'(x)| \leq \frac{V_n(x)}{\rho_n}, \quad 0 < V_n''(x) \leq \frac{2V_n(x)}{\rho_n^2}, \quad (2.22)$$

for all $x \in g_n$. Define the numbers $M_n = \max_{x \in g_n} V_n(x)$, $M_n' = \max_{x \in g_n} V_n'(x)$. Let $Q_p = \frac{1}{\pi} \int_{\mathbb{R}} x^{2p} v(x) dx$, $p \geq 0$. We prove Theorem 1.5 for more general case.

Lemma 2.4. *Let $u_* = \inf_{n \in \mathbb{Z}} (u_{n+1} - u_n) > 0$ and let $Q_0 < \infty$. Then*

$$M_n \leq \frac{2}{\pi \rho_n} \sup_{s \in \mathbb{Z}} h_s \leq \frac{\sqrt{8Q_0}}{\pi \rho_n}, \quad M_n' \leq \frac{M_n}{\rho_n}, \quad M_n'' \leq \frac{M_n}{\rho_n^2}, \quad (2.23)$$

$$\sum M_n \leq \frac{\pi^2}{3\rho^2} Q_0, \quad (2.24)$$

$$\rho = \inf \rho_n \geq \frac{2u_*}{5\pi} \exp\left(-\frac{5\pi}{2u_*} \sup_{s \in \mathbb{Z}} h_s\right). \quad (2.25)$$

If additionally $Q_2 < \infty$, then

$$V_n(x) \leq \frac{2}{|x|\sqrt{|z_n^+ z_n^-|}} \left(Q_0 + \frac{Q_2}{\rho_n^2}\right), \quad x \in \bar{g}_n. \quad (2.26)$$

Proof. Let $h_+ = \sup h_n$. We have for $x \in g_n$

$$V_n(x) = \frac{1}{\pi} \int_{g \setminus g_n} \frac{v(t)dt}{|t-x|v_n(t)} \leq \frac{h_+}{\pi} \int_{g \setminus g_n} \frac{dt}{|t-x|v_n(t)} \leq \frac{h_+}{\pi} \int_{|s|>\rho_n} \frac{ds}{|s|^2} \leq \frac{2h_+}{\pi\rho_n}$$

and using $h_+^2 \leq 2Q_0$ (see [K5]) we obtain the first estimate in (2.23). The two last estimates follow from (2.22).

Let $M_n = V_n(x_n)$ for some $x_n \in g_n$. Using $|t - x_n| \geq \rho|m - n|$ for any $t \in g_m$, we obtain

$$\sum M_n \leq \sum_n \sum_{m \neq n} \frac{1}{\pi} \int_{g_m} \frac{v(t)dt}{|t - x_n|v_n(t)} \leq \sum_n \sum_{m \neq n} \frac{1}{\pi} \int_{g_m} \frac{v(t)dt}{\rho^2|n - m|^2} = \frac{Q_0 C}{\rho^2},$$

where $C = 2 \sum_{n \geq 1} n^{-2} = 2 \frac{\pi^2}{6} = \frac{\pi^2}{3}$. Estimate (2.25) was proved in [KK4].

Using the simple estimate $\frac{1}{|t-z|} \leq \frac{1}{|z|} \left(1 + \frac{|t|}{\rho_n}\right)$ for all $z \in \bar{g}_n, t \in g \setminus g_n$, we obtain

$$\begin{aligned} V_n(x) &= \frac{1}{\pi} \int_{g \setminus g_n} \frac{v(t)dt}{|t-x|v_n(t)} \leq \frac{1}{\pi} \int_{g \setminus g_n} \left(1 + \frac{|t|}{\rho_n}\right)^2 \frac{v(t)dt}{|x|\sqrt{|z_n^+ z_n^-|}} \\ &\leq \frac{2}{\pi|x|\sqrt{|z_n^+ z_n^-|}} \int_{\mathbb{R}} \left(1 + \frac{t^2}{\rho_n^2}\right) v(t)dt \leq \frac{2}{|x|\sqrt{|z_n^+ z_n^-|}} \left(Q_0 + \frac{Q_2}{\rho_n^2}\right), \end{aligned}$$

which yields (2.26). ■

3 Estimates for the Hill and the Dirac operators

Firstly, we will apply the results from Section 2 for the case the Dirac operator T_D .

Proof of Theorem 1.3. Estimate (1.17), (1.18), (1.19), (1.20), (1.21) follows from (2.4), (2.6), (2.17), (2.18), (2.19) respectively. ■

Proof of Theorem 1.4. Estimate (1.23), (1.24), (1.26), (1.27) follows from (2.10), (2.12), (2.13), (2.14). ■

Secondly we will apply the results from Section 2 for the case the Hill operator $T = -\frac{d^2}{dt^2} + (q_0 + q'(t))$ in $L^2(\mathbb{R})$ with the 1-periodic real potential $q \in L^2(0, 1)$. Recall that $\lambda_0^+ = 0$ for the Hill operator. In this case we have estimates

$$\frac{|\gamma_n|}{2\sqrt{\lambda_n^+}} \leq |g_n| = \frac{|\gamma_n|}{z_n^+ + z_n^-} \leq \frac{|\gamma_n|}{2n\rho}, \quad m_n^\pm = 2z_n^\pm \mu_n^\pm, \quad n \geq 1. \quad (3.1)$$

Proof of Theorem 1.1. Recall $\lambda_n^0 = \frac{\lambda_n^+ + \lambda_n^-}{2}$ and $\lambda_n = z_n^2$. Using the identity

$$4(z_n^0)^2 = (z_n^- + z_n^+)^2 = \lambda_n^- + \lambda_n^+ + 2z_n^- z_n^+ = 4\lambda_n^0 - |g_n|^2,$$

and the estimate (2.4) we obtain

$$|\lambda_n^0 - \frac{|g_n|^2}{4} - \lambda_n| = |(z_n^0)^2 - \lambda_n| = |z_n^0 + z_n||z_n^0 - z_n| \leq |z_n^0 + z_n| \frac{|g_n|^2}{4} |V'_n(z_n)| \leq \frac{3|\gamma_n|^2}{8z_n^0} |V'_n(z_n)|,$$

which yields (1.4). Recall $2z_n^\pm \mu_n^\pm = m_n^\pm$. Using (2.6) we obtain (1.5).

Using (2.17) and $|g_n||m_n^\pm| \leq 2|\gamma_n||\mu_n^\pm|$ we obtain (1.6).

Using (2.18), (2.9) and $2z_n^\pm \mu_n^\pm = m_n^\pm$ we obtain

$$|\mu_n^+ + \mu_n^-| = \frac{|z_n^0(m_n^+ + m_n^-) + c_n(m_n^- - m_n^+)|}{2z_n^+ z_n^-} \leq \frac{z_n^0 |g_n|^2}{2z_n^+ z_n^-} (1 + M_n) M'_n + \frac{|g_n|^2}{4z_n^+ z_n^-} (1 + M_n),$$

which yields (1.7). Using $\mu_n = \frac{m_n}{2z_n^0} = \frac{2h_n^2}{|\gamma_n|}$ we obtain $\mu_n^\pm \mp \mu_n = \frac{\mp |g_n| m_n^\pm}{4z_n^0 z_n^\pm} + \frac{m_n^\pm - m_n}{2z_n^0}$. The estimate (2.19) yields (note that $|g_n||m_n^\pm| \leq 2|\gamma_n||\mu_n^\pm|$)

$$|m_n^\pm - m_n| \leq |g_n|^2 (1 + M_n) M'_n \left(1 + \frac{M'_n}{8} \sqrt{2|g_n||m_n^\pm|} \right)$$

and (2.9) gives $|m_n^\pm| \leq |g_n|(1 + M_n)^2/2$. The two last estimates imply (1.8). ■

Proof of Theorem 1.2. Using (2.11) we obtain

$$A_n = \frac{4}{\pi} \int_{g_n} x v(x) dx = 4z_n^0 a_n + A_{n0}, \quad A_{n0} = \frac{4}{\pi} \int_{g_n} (x - z_n^0) v(x) dx, \quad |A_{n0}| \leq \frac{|g_n|^4}{2^5} M'_n. \quad (3.2)$$

Estimate (2.10) and $2z_n^0 |g_n| = |\gamma_n|$ imply

$$|4z_n^0 a_n - (|\gamma_n|^2/8z_n^0)(1 + V_n(z_n^0))| \leq z_n^0 \frac{|g_n|^4}{2^5} M''_n, \quad (3.3)$$

which together with (3.2) yields (1.10). Estimate (2.12) implies

$$|4z_n^0 a_n - \frac{|\gamma_n| h_n}{2}| \leq z_n^0 \frac{|g_n|^4}{2^5} (M''_n + 6M'_n{}^2),$$

which together with (3.2) yields (1.11). Estimate (2.13) implies

$$|4z_n^0 a_n - \frac{|\gamma_n|}{2} |c_n m_n^\pm|^{\frac{1}{2}}| \leq 2z_n^0 c_n^3 (M'_n + c_n M''_n)$$

which together with (3.2) yields (1.12). Using and $\int_{g_n} (x - z_n^0) v_n(x) dx = 0$, we obtain

$$\beta_n^2 = A_n = \frac{4}{\pi} \int_{g_n} x v(x) dx \geq \frac{4}{\pi} \int_{g_n} x v_n(x) dx = \frac{4z_n^0}{\pi} \int_{g_n} v_n(x) dx = \frac{z_n^0 |g_n|^2}{2} \quad (3.4)$$

Recall that (1.10) yields $\left| A_n - |\gamma_n|^2 \frac{(1+V_n(z_n^0))}{8z_n^0} \right| \leq \frac{|g_n|^4}{2^5} (z_n^0 M_n'' + M_n')$, which gives

$$\left| \beta_n - |\gamma_n| \sqrt{\frac{(1+V_n(z_n^0))}{8z_n^0}} \right| \leq \frac{|g_n|^4}{2^5(\beta_n + |g_n| \sqrt{z_n^0/2})} (z_n^0 M_n'' + M_n'),$$

and (3.4) yields (1.13). Using (1.11), (1.12) and the similar arguments we obtain (1.14), (1.15). $\blacksquare$

Proof of Corollary 1.6. Using (1.31), (1.32) we obtain

$$\sum_{p \in \mathbb{Z}} M_p'' \leq \frac{1}{\rho^2} \sum_{p \in \mathbb{Z}} M_p \leq \frac{\pi^2 Q_0}{3\rho^4}, \quad \sum_{p \in \mathbb{Z}} M_p'^2 \leq \frac{\sqrt{8Q_0}}{\pi\rho} \frac{\pi^2 Q_0}{3\rho^4},$$

which yields

$$\sum_{p \in \mathbb{Z}} (M_p'' + 6M_p'^2) \leq \frac{\pi^2 Q_0}{3\rho^4} \left(1 + 6 \frac{\sqrt{8Q_0}}{\pi\rho} \right). \quad (3.5)$$

Using (1.24) and (3.5) we obtain

$$\begin{aligned} \sum_{n \in \mathbb{Z}} \left| a_n - \frac{|g_n| h_n}{4} \right|^{\frac{1}{3}} &\leq \sum_{n \in \mathbb{Z}} \frac{|g_n|^{\frac{4}{3}}}{2^{\frac{7}{3}}} (M_n'' + 6M_n'^2)^{\frac{1}{3}} \leq 2^{\frac{-7}{3}} \left(\sum_{n \in \mathbb{Z}} |g_n|^2 \right)^{\frac{2}{3}} \left(\sum_{p \in \mathbb{Z}} (M_p'' + 6M_p'^2) \right)^{\frac{1}{3}} \\ &\leq 2^{\frac{-7}{3}} \left(\frac{Q_0}{8} \right)^{\frac{2}{3}} \left(\frac{\pi^2 Q_0}{3\rho^4} \right)^{\frac{1}{3}} \left(1 + \frac{\sqrt{8Q_0}}{\pi\rho} \right)^{\frac{1}{3}}, \end{aligned}$$

which yields (1.35). Using (1.24) and (3.5) we obtain

$$\sum_{p \in \mathbb{Z}} \left| a_n - \frac{|g_n| h_n}{4} \right| \leq \sum_{p \in \mathbb{Z}} \frac{|g_n|^4}{2^7} (M_n'' + 6M_n'^2) \leq \frac{g_+^4}{2^7} \frac{\pi^2 Q_0}{3\rho^4} \left(1 + 6 \frac{\sqrt{8Q_0}}{\pi\rho} \right), \quad g_+ = \sup |g_n|.$$

$\blacksquare$

References

- [FM] Flaschka, H.; McLaughlin, D. Canonically conjugate variables for the Korteweg- de Vries equation and the Toda lattice with periodic boundary conditions. Prog. of Theor. Phys. 55(1976), 438-456,
- [GT] Garnett, J.; Trubowitz, E. Gaps and bands of one dimensional periodic Schrödinger operators. Comment. Math. Helv. 59(1984), 258-312,
- [G] Golusin, G. Geometric theory of functions of a complex variable, Amr. Math. Soc., Providence, RI, 1969,
- [J] Jenkins, A. Univalent functions and conformal mapping. Berlin, Göttingen, Heidelberg: Springer, 1958,

- [JM] Johnson, R.; Moser, J. The rotation number for almost periodic potentials. *Commun. Math. Phys.* 84(1982), 403-430,
- [KM] Kappeler, T.; Möhr, C. Estimates for periodic and Dirichlet eigenvalues of the Schrödinger operator with singular potentials. *J. Funct. Anal.* 186 (2001), no. 1, 62–91,
- [KK1] Kargaev, P.; Korotyaev, E. Effective masses and conformal mappings. *Commun. Math. Phys.* 169(1995), 597-625,
- [KK2] Kargaev, P.; Korotyaev, E. The inverse problem for the Hill operator, a direct method. *Invent. Math.* 129(1997), 567-593,
- [KK3] Kargaev, P.; Korotyaev, E. Inverse electrostatic problems on plane, in preparation,
- [KK4] Kargaev, P.; Korotyaev, E. A priori estimates for conformal mappings on complex plane with parallel slits, preprint 2006,
- [K1] Korotyaev, E. Metric properties of conformal mappings on the complex plane with parallel slits. *Inter. Math. Reseach. Notices.* 10(1996), 493-503
- [K2] Korotyaev, E. Estimates for the Hill operator.I, *Journal Diff. Eq.* 162(2000), 1-26,
- [K3] Korotyaev, E. Estimate for the Hill operator.II, *J. Diff. Equations* 223 (2006), 229–260,
- [K4] Korotyaev, E. The estimates of periodic potentials in terms of effective masses. *Commun. Math. Phys.* 183(1997), 383-400,
- [K5] Korotyaev, E. Estimate of periodic potentials in terms of gap lengths. *Commun. Math. Phys.* 197(1998), 521-526,
- [K6] Korotyaev, E. Inverse oroblem and estimates for periodic Zakharov-Shabat systems, *J. Reiner Angew. Math.* 583(2005), 87-115,
- [K7] Korotyaev, E. Characterization of the spectrum of Schrödinger operators with periodic distributions. *Int. Math. Res. Not.* (2003) no. 37, 2019–2031,
- [K8] Korotyaev, E. Inverse problem for periodic "weighted" operators. *J. Funct. Anal.* 170 (2000), 188–218,
- [K9] Korotyaev, E. Periodic "weighted" operators. *J. Differential Equations* 189 (2003), no. 2, 461–486,
- [Kr] Krein M. On the characteristic function $A(\lambda)$ of a linear canonical system differential equation of the second order with periodic coefficients (Russian) , *Prikl. Mat. Meh.* 21(1957), 320-329,
- [Le] Levin, B. Majorants in the class of subharmonic functions.1-3. *Theory of functions, functional analysis and their applications.* 51(1989), 3-17 ; 52(1989), 3-33 . Russian,
- [LS] Levitan, B.; Sargsjan, I. Sturm-Liouville and Dirac operators. Translated from the Russian. *Mathematics and its Applications (Soviet Series)*, 59. Kluwer Academic Publishers Group, Dordericht, 1991,

- [MO1] Marchenko, V.; Ostrovski, I. A characterization of the spectrum of the Hill operator. Math. USSR Sb. 26(1975), 493-554
- [MO2] Marchenko, V.; Ostrovski, I. Approximation of periodic by finite-zone potentials. Selecta Math. Sovietica. 6(1987), 101-136
- [NS] Neĭman-zade, M. I.; Shkalikov, A. A. Schrödinger operators with singular potentials from spaces of multipliers. (Russian) Mat. Zametki 66 (1999), no. 5, 723–733,
- [SS] Savchuk, A. M.; Shkalikov, A. A. Sturm-Liouville operators with singular potentials. (Russian) Mat. Zametki 66 (1999), no. 6, 897–912.